

Normal and Standard Normal Distribution

by James R. Garven*
December 21, 2016

- A continuous random variable x has a *normal distribution* if its probability density function is

$$f(x) = \frac{e^{-.5((x-\mu_x)/\sigma_x)^2}}{\sigma_x\sqrt{2\pi}},$$


where $\sigma_x > 0$, $-\infty < \mu_x < \infty$, $-\infty < x < \infty$, and $\int_{-\infty}^{\infty} f(x)dx = 1.0$.

- The normal probability density function has two parameters: μ_x (mean, or expected value) and σ_x (standard deviation).

- Next, we define the *standard normal distribution*. This involves transforming the normal random variable x into a standard normal random variable $z = (x-\mu_x)/\sigma_x$. Applying the expected value operator to both sides of this equation, $E(z) = (E(x)-\mu_x)/\sigma_x = 0$ (since $E(x) = \mu_x$). Furthermore, $\sigma_z^2 = E[(z - E(z))^2] = E(z^2)$ (since $E(z) = 0$). Therefore, $\sigma_z^2 = E(z^2) = E\left(\left\{\frac{x - \mu_x}{\sigma_x}\right\}^2\right) = (1/\sigma_x^2) E[(x - \mu_x)^2] = (\sigma_x^2/\sigma_x^2) = 1$, which implies that $\sigma_z = 1$. Thus,

the *standard normal density function* is $f(z) = \frac{e^{-z^2/2}}{\sqrt{2\pi}}$.


- Here is a graph of the standard normal density function, $f(z)$:



*James R. Garven is the Frank S. Groner Memorial Chair in Finance and Professor of Finance & Insurance at Baylor University (Address: Foster 320.39, One Bear Place #98004, Waco, TX 76798, telephone: 254-307-1317, e-mail: James_Garven@baylor.edu).

Note that $f(z)$ is symmetric; half of its area is to the left of 0, and the other half is located to the right. Furthermore, although z is unbounded from below and above, nearly 100% of its area occurs within plus and minus 4 standard deviations from its mean (recall that $\sigma_z = 1$).

- Next, we calculate and graph the cumulative distribution function for z , $F(z) = \int_{-\infty}^z f(z)dz$:



Note that to the left of z 's mean of 0, $F(z)$ is convex (increasing at an increasing rate), whereas to the right of 0, $F(z)$ is concave (increasing at a decreasing rate). As z becomes a "large" positive number, $F(z)$ approaches a value of 1. Also, the slope of $F(z)$ is equal to $f(z)$ for all values of z .

- Calculating $F(z)$: Thankfully, we don't have to literally integrate the standard normal density function in order to calculate probabilities. Excel has a built-in function called NORMSDIST; when you use the command =NORMSDIST(z), Excel returns the probability. For example, if you type =NORMSDIST(0), Excel performs the following calculation: $F(0) = \int_{-\infty}^0 f(z)dz = 50\%$. If you were interested in knowing what the probability is of being within plus or minus 1 standard deviation from the mean, then by typing =NORMSDIST(1)-NORMSDIST(-1), Excel performs the following calculation: $F(1) - F(-1) = \int_{-1}^1 f(z)dz = 68.27\%$, and so forth. You can also perform these same calculations by referencing a "z" table (e.g., from the home page of the course website, click on "[Formula Sheets](#)", then "[Standard Normal Distribution Function \("z"\) Table](#)").